



Journal of Environmental Science and Technology

ISSN 1994-7887

science
alert

ANSI*net*
an open access publisher
<http://ansinet.com>

Metal Hyperaccumulation in Plants: A Review Focusing on Phytoremediation Technology

Hemen Sarma

Centre for Advanced Studies in Botany, School of Life Science, North-Eastern Hill University, Shillong-7930 22, India

ABSTRACT

Metal hyperaccumulation is a characteristic present in over 500 plant species and approximately in 0.2% of all angiosperms. Hyperaccumulators are model plants for phytoremediation as they are tolerant to heavy metals. Metals hyperaccumulation and tolerance are genetically inherited traits. Plants possess a range of potential cellular mechanisms that may be involved in the detoxification of heavy metals and thus tolerance to metal stress. Recent discovery have given first insights into the molecular basis of metal hyperaccumulation and metal hypertolerance in some plants. However, the ecological and biological significance of hyperaccumulation is not clear yet. The recent progress of molecular techniques has helped to improve the performance of phytoremediation technology as well as plant adaptation to extreme metallic environments. The knowledge of metal hyperaccumulation physiology has recently developed as a result of the advancement of molecular biology. Molecular technique help to understand the gene regulations system and plant metal homeostasis. This study reviews the recent advances of phytoremediation technology using hyperaccumulator plants addressing both potential and limitations, physiological and molecular aspects and provides a broad overview of most important genes which have been correlated to metals hyperaccumulation and tolerance, evidence of the effect of heavy metal on biomass productions, plant biochemical, antioxidant defence system and discusses the prospects of transgenic plants in phytoremediation of heavy metals.

Key words: Antioxidant, biomass, chlorophyll, heavy metals, metal hyperaccumulation, molecular biology, physiology, phytoremediation, transgenic plants

INTRODUCTION

In recent years, public concerns relating to ecological threats caused by heavy metal (HM) have led to intensive research of new economical plants based remediation technologies. Conventional methods used for reclamation of contaminated soils, namely chemical, physical and microbiological methods, are costly to install and operate (Danh *et al.*, 2009). The rapid increase in population coupled with fast industrialization growth causes serious environmental problems, including the production and release of considerable amounts of toxic waste materials into environment (Zhuang *et al.*, 2007). Over recent decades, the annual worldwide release of heavy metals reached 22,000 t (metric ton) for cadmium, 939,000 t for copper, 783,000 t for lead and 1,350,000 t for zinc (Singh *et al.*, 2003). Vanadium released from fertilizer effluents usually ranges between 16.58-18.44 mg L⁻¹ (Sarma, 2008). However, the combustion of coal and petroleum-based products is considered as the primary source of vanadium in the environment (Adriano, 1986). According to an estimate, in US approximately 2125 ton of elemental vanadium has escaped in to the environment annually (Kaplan *et al.*, 1990).

In an ecological research, any metal or metalloid that causes environmental problem which cannot be biologically degraded should be considered as a heavy metal. Heavy metals are natural components of the Earth's crust, but in many ecosystems the concentration of several HMs has reached toxic levels due to consequence of anthropogenic activities. Fifty three elements fall into the category of heavy metal till date and defined as the group of elements whose densities are higher than 5 g cm^{-3} and recognized as ubiquitous environmental contaminants in industrialized societies (Padmavathiamma and Loretta, 2007). The adaptative responses of plants to HM contaminated environments are efficient processes that include many physiological, molecular, genetic and ecological traits. These traits give certain species the ability to survive or hyperaccumulate the toxic metals. Soil pollution caused by metals is some what different from air or water pollution, because heavy metals persist in soil much longer periods of time than in other compartments of the biosphere (Lasat, 2002). Out of the 92 known elements present on earth crust, only 17 are known to be essential to all living plants for normal growth and 12 are potentially beneficial required only in trace amounts includes silver (Ag), cerium (Ce), chromium (Cr), fluoride (F), iodine (I), lanthanum (La), rubidium (Rb), tin (Sn), serium (Sr), titanium (Ti), vanadium (V) and tungsten (W) (Pilon-Smits *et al.*, 2009).

Mechanisms of carcinogenicity has been discussed for metals and classified arsenic, antimony, beryllium, cadmium, chromium, cobalt, lead, nickel and vanadium are some metals and metalloids that are carcinogenic to humans or considered to be carcinogenic to humans by International Agency for Research on Cancer (Beyersmann and Hartwig, 2008). Toxic heavy metals cause DNA damage and their carcinogenic effects in animals and humans are probably caused by their mutagenic ability (Knasmuller *et al.*, 1998; Baudouin *et al.*, 2002). Exposure to high levels of these metals has been linked to adverse effects on human health and wildlife.

The hyperaccumulation of heavy metals in some plants has been recorded by many researchers during last few decades (Barman *et al.*, 2000) and this has emphasized the importance of further advanced research in molecular basis of phytoremediation technology. The hyperaccumulation of heavy metals is depends on the plant species, soil condition (pH, organic matter content, cation exchange capacity etc.) and types of heavy metal (Barman *et al.*, 2001; Spinoza-Quinones *et al.*, 2005; Xian and Shokohifard, 1989; Otte *et al.*, 1993). In metal biology, it is experimentally proved that even some metals that are essential for the normal plants growth (such as iron and copper) may become toxic, depending on the oxidation state, complex form, dose and mode of exposure (Beyersmann and Hartwig, 2008).

The removal and recovery of heavy metals from contaminated media are of great importance in terms of protection of the environment as well as abatement of heavy metal toxicity (Kim *et al.*, 2004). Metals and their compounds are accumulating up to harmful levels, unlike organic waste by some living plants. The level of toxic metals (Pb, Cr, Hg etc.) can be reduced from contaminated media by a number of aquatic plants taken up by the roots system and transported to the stems and leaves without showing toxicity syndrome has confirmed by many studies (Rai *et al.*, 1995; Cardwell *et al.*, 2002). This uptake of metals largely depends on the type and chemical speciation of metal and habitat characteristics of plants, i.e., terrestrial, aquatic etc. Hence, plant selection is important for remediation of contaminated site.

Remediation of metal-contaminated soils became a goal for many research laboratories in the world. The use of plants in designing low cost treatment system is still a challenge in environmental managements. Phytoremediation, a new green technology has been inventoried, in which some metal hyperaccumulator plants were utilized to decontaminate soil, water and ambient

environment (Prasad, 2003) and it is growing without symptoms of toxicity (Reeves and Brooks, 1983; Baker and Brooks, 1989; Baker *et al.*, 1991).

Hyperaccumulator plants represent a resource for remediation of metal polluted site, as they are able to extract wide range of metals and to concentrate them in their upper parts with the character of metal tolerance. In some plant species, the concentrations accumulated in aboveground biomass of metals or metalloids are more than one and up to four, orders of magnitude higher than in other adjacent plants (Baker and Brooks, 1989; Reeves and Baker, 2000). This unique extent of accumulation of heavy metals, have been reported in a total of approximately 500 plant species to date (Kramer, 2010). The first field trial on Zn and Cd phytoextraction was conducted by Baker *et al.* (1991) and now a days, this technology is receiving considerable attention for clean up of soil contaminated with heavy metals (USEPA, 2000). Several comprehensive reviews have been published, summarizing many important aspects of this novel green technology (Padmavathamma and Loretta, 2007; Kramer, 2010). The present reviews give general guidance of research trends in last few decades, focusing the physiological and molecular basis of phytoremediation. For this review potential publications were searched using Biological Abstracts (BIOSIS), Biological Sciences (CSA), Biosis Previews (BIOSIS), CAB Abstracts (CAB International), Plant Science (CSA) and Web of Science (ISI) data bases; the last round of the search was conducted in June 2010. We used key words that referred to phytoremediation for searching (different combinations of keywords: metals *, plants, hyperaccumulator*, physiology*, Phytochelatins biochemical*). These keywords produced a reasonable output that was then sorted manually; However, the list is not exclusive. In addition, we checked the reference sections of all discovered publications for references to other potentially suitable studies as a cross reference. This review specially focused update information so far available and provides a critical overview of the present state of the art.

Definitions: The term phytoremediation (phyto meaning plant and the Latin suffix remedium meaning to clean or restore) refers to a diverse collection of plant-based technologies that use either naturally occurring, or genetically engineered, plants to clean contaminated environments (Flathman and Lanza, 1998). Phytoremediation is clean, simple, cost effective, non-environmentally disruptive (Wei *et al.*, 2004) green technology and most importantly, its by-products can find a range of other uses (Truong, 1999, 2003).

Phytoremediation is an eco friendly approach for remediation of contaminated soil and water using plants comprised of two components, one by the root colonizing microbes and the other by plants themselves, which accumulates the toxic compounds to further non toxic metabolites. Various compounds viz., organic synthetic compounds, xenobiotics, pesticides, hydrocarbon and heavy metals, are among the contaminants that can be effectively remediated by plants (Suresh and Ravishankar, 2004).

TYPES OF PHYTOREMEDIATION TECHNOLOGY

The four different plant-based technologies of phytoremediation, each having a different mechanism of action for remediating metal-polluted soil, sediment or water:

- **Phytoextraction:** Plants absorb metals from soil through the root system and translocate them to harvestable shoots where they accumulate. Hyperaccumulators mostly used this process to extract metals from the contaminated site. The recoveries of the extracted metals are also possible through harvesting the plants appropriately

Table 1: Cost of different remediation technologies (Glass, 1999)

Treatments	Cost (US\$ ton ⁻¹)
Vitrification	75-425
Land filling	100-500
Chemical treatments	100-500
Electrokinetics	20-200
Phytoextraction	5-40

- **Phytovolatilization:** Plants used to extract certain metals from soil and then release them into the atmosphere by volatilization
- **Phytostabilization:** In this process, the plant roots and microbial interactions can immobilized organic and some inorganic contaminants by binding them to soil particles and as a result reduce migration of contaminants to grown water
- **Phytofiltration:** Phytofiltration is the use of plants roots (rhizofiltration) or seedlings (blastofiltration) to absorb or adsorb pollutants, mainly metals, from water and aqueous waste Streams (Prasad and Freitas, 2003)

Limitations: The application of phytoremediation for pollution control has several limitations that require further intensive research on plants and site-specific soil conditions (Danh *et al.*, 2009). It is generally slower than most other treatment viz., chemical, physical and microbiological plants with low biomass yields and reduced root systems do not support efficient phytoremediation and most likely do not prevent the leaching of contaminants into aquatic system. Environmental conditions also determine the efficiency of phytoremediation as the survival and growth of plants are adversely affected by extreme environmental conditions, toxicity and the general conditions of soil in contaminated lands (Danh *et al.*, 2009). In phytoremediation technology, multiple metals contaminated soil and water requires specific metal hyperaccumulator species and therefore requires a wide range of research prior to the applications. Though the phytoremediation is cost effective (Table 1), environment friendly, ability to reclaim heavy metals contaminated site, several limitations also create trouble in implementing the strategy, e.g., metal must be in bio-available form to plants; if metals is tightly bound to the organic portions of the soil, some time it may not be available to plants. Furthermore, if the metals are water soluble, in nature it will pass by the root system without accumulation.

The phytoremediation of mixed heavy metals contaminated soil have conformant with some problem e.g., The cadmium/zinc model hyperaccumulator *Thlaspi caerulescens* is sensitive toward copper (Cu) toxicity, which is a problem in remediation of Cd/Zn from soils in the presence of Cu by application of this species. In *T. caerulescens* Cu induced inhibition of photosynthesis followed the sun reaction type of damage, with inhibition of the photosystem II reaction center charge separation and the water-splitting complex (Mijovilovich *et al.*, 2009). Despite some limitations, present day phytoremediation technology are using worldwide and various research laboratories are at present engaged to overcome the limitations.

PLANTS SELECTIONS CRITERIA FOR PHYTOREMEDIATION

Plant species selection is a critical management decision for phytoremediation. Grasses are thought to be excellent candidates, because their fibrous rooting systems can stabilize soil and provide a large surface area for root-soil contact (Kulakow *et al.*, 2000). The selection of plants is

possibly the single most important factor for fruitful phytoremediation strategy. The application of indigenous plant species for phytoremediation is often favoured as it requires less management and acclimatizes successfully in native climate conditions and seasonal cycle. However, some exotic plant species may perform better in remediation of specific metals and can be safely used where the possibility of invasive behavior has been eliminated (USEPA, 2000). Some important criteria in selecting plant species for phytoremediation are:

- The levels of tolerance with respect to metal known to exist at the site
- The level of adequate accumulation, translocation and uptake potential of metals
- High growth rate and biomass yield
- Tolerance to water logging and extreme drought conditions
- Availability, habitat preference e.g., terrestrial, aquatic, semi-aquatic etc.
- Tolerance to high pH and salinity
- Root characteristic and depth of the root zone

METAL HYPERACCUMULATORS PLANTS/FAMILIES

Over 500 plant species comprising of 101 families have been reported, including members of the Asteraceae, Brassicaceae, Caryophyllaceae, Cyperaceae, Cunouniaceae, Fabaceae, Flacourtiaceae, Lamiaceae, Poaceae, Violaceae and Euphobiaceae. Metal hyperaccumulation occurs in approximately 0.2% of all angiosperms and is particularly well represented in the Brassicaceae (Kramer, 2010). Recently Environment Canada has released a database Phytorem which compiled a worldwide inventory of more than 750 terrestrial and aquatic plants with potential value for phytoremediation. The study of Co accumulations in 670 species of terrestrial plants showed that leaf Co concentration was in general less than 0.2 ppm, with the exception of Ericales, Euasterids and Asparagales, where 0.3-0.5 ppm of Co was measured (Watanabe *et al.*, 2007). A list of 26 Co hyperaccumulators species belonging to the families of Lamiaceae, Scrophulariaceae, Asteraceae and Fabaceae has been reported (Baker *et al.*, 2000). Zn hyperaccumulation was first discovered in 1865 in *Noccaea caerulescens* (syn. *Thlaspi caerulescens*) of Brassicaceae (Reeves and Baker, 2000) and Ni hyperaccumulation, which was first reported in 1948 in *Alyssum bertolonii* of Brassicaceae (Kramer, 2010). A member of Crassulaceae *Sedum alfredii* is the only known Cd hyperaccumulator outside the Brassicaceae (Deng *et al.*, 2007). The hyperaccumulation of arsenic (As) has been discovered in only two species of the Brassicaceae (Karimi *et al.*, 2009) apart from a number of pteridophytes (Ma *et al.*, 2001).

METAL HYPERACCUMULATOR MODEL PLANTS

Mixed pollution with heavy metals is characteristic for many spill areas and industrial effluents dumping sites. The danger for the environment and human health from such sites is large and sustainable remediation strategies are urgently needed. Phytoremediation seems to be cheap and environmentally sound option for reclaim the hazardous toxic metals and metalloids. The most important challenge is how to improve the efficiency of phytoremediation by increasing the accumulation of metals in plants, or by improving key plant biological traits that should enhance metal uptake (Wu and Tang, 2009). The heavy metals accumulation on several plants species have been focused viz., *Thlaspi* sp. (Baker *et al.*, 1994), *Brassica* sp. (Blaylock *et al.*, 1997; Huang *et al.*, 1997) and *Alyssum* (Kramer *et al.*, 1996). Heavy metal accumulators is increasing steadily (currently more than 500 plant species) and some are presented in Table 2.

Table 2: Examples of some plants and metals they can remediate

Species	Metal	References
<i>Alyssum wulfenianum</i>	Ni	Reeves and Brooks (1983)
<i>Azolla pinnata</i> , <i>lemna minor</i>	Cu, Cr	Jain <i>et al.</i> (1990)
<i>Brassica Juncea</i>	Cu, Ni	Ebbs and Kochian (1997)
<i>Arobiodopsis hallerii</i>	Cd	Kupper <i>et al.</i> (2000)
<i>Pteris vittata</i>	Cu, Ni, Zn	Ma <i>et al.</i> (2001)
<i>Psychotria douarrei</i>	Ni	Davis <i>et al.</i> (2001)
<i>Pelargonium</i> sp.	Cd	Dan <i>et al.</i> (2002)
<i>Thlaspi caerulescens</i>	Zn, Cd and Ni	Assuncao <i>et al.</i> (2003)
<i>Arabidopsis halleri</i>	Cd	Berts and Meerts <i>et al.</i> (2003)
<i>Amanita muscaria</i>	Hg	Falandysz <i>et al.</i> (2003)
<i>Arabis gemmifera</i>	Cd and Zn	Kubota and Takenka (2003)
<i>Pistia stratiotes</i>	Ag, Cd, Cr, Cu, Hg, Ni, Pb and Zn	Odjegba and Fasidi (2004)
<i>Piptathertan miliacetall</i>	Pb	Garcia <i>et al.</i> (2004)
Spartina plants	Hg	Tian <i>et al.</i> (2004)
<i>Astragalus bisulcatus</i> , <i>Brassica Juncea</i>	Selenium	Ellis <i>et al.</i> (2004)
<i>Sedum alfredii</i>	Cd	Xiong <i>et al.</i> (2004)
<i>H. annuus</i>	Pb	Boonyapookana <i>et al.</i> (2005)
<i>H. indicus</i>	Pb	Chandra Sekhar <i>et al.</i> (2005)
<i>Sesbania drummondii</i>	Pb	Sharma <i>et al.</i> (2004)
<i>Lemna gibba</i>	As	Mkandvire and Dude (2005)
<i>Pteris vittata</i>	AS	Dong (2005)
<i>Sedum alfredii</i>	Pb/Zn	Sun <i>et al.</i> (2005)
<i>Thlaspi caerulescens</i>	Zn, Pb, Zn and Cd	Banasova and Horak (2008)
<i>Chengiopanax sciadophylloides</i>	Mn	Mizuno <i>et al.</i> (2008)
<i>Tamarix smyrnensis</i>	Cd	Manousaki <i>et al.</i> (2008)
<i>P. griffithii</i>	Cd/Zn	Hu <i>et al.</i> (2009)
<i>Brassica napus</i>	Cd	Selvam and Wong (2008)
<i>Arabidopsis thaliana</i>	Zn and Cd	Saraswat and Rai (2009)
<i>Crotalaria juncea</i>	Ni and Cr	Saraswat and Rai (2009)
<i>C. dactylon</i>	Ni and Cr	Saraswat and Rai (2009)
<i>Rorippa globosa</i>	Cd	Sun <i>et al.</i> (2010)

Some species are highly metal specific, have a small biomass, slow growth habit and require careful management for multiplications (Gleba *et al.*, 1999) which are not suitable for commercial applications. Thus identification of novel plant species with high biomass yield coupled with ability to tolerate and accumulate multiple metals has become an important aspect of phytoremediation research. A multiple metal hyperaccumulator scented geranium (*Pelargonium* sp.) has been recently discovered. This plant has an ability to tolerate and accumulate multiple metals (Cd, Ni and Pb) and maintain normal metabolic processes (Dan *et al.*, 2000; KrishnaRaj *et al.*, 2000). Another potential Pb hyperaccumulator perennial shrub *Sesbania drummondii* with high biomass yield has been discovered from southern coastal areas of the United State (Sahi *et al.*, 2002) was given as evidence for model plants.

In the last few decades many scientists in different parts of the worlds has worked out the metals bioaccumulation potential of various species and some are presented in Table 3 derived through meta-analysis.

Works on physiological aspects of *Vetiveria zizanioides* has provided an extensive research foundation which makes this plant a good candidate for a wide range of phytoremediation purposes

Table 3: Examples of some metal hyperaccumulator and their bioaccumulation potential

Plants	Metal	Bioaccumulation	References
<i>A. racemosus</i>	Se	14,900 mg kg ⁻¹	Beath <i>et al.</i> (1937)
<i>Sebertia acuminata</i>	Ni	25% by wt. dried sap	Jaffre <i>et al.</i> (1976)
<i>Ipomea alpine</i>	Cu	12,300 mg kg ⁻¹	Baker and Walker (1989)
<i>Berkheya coddii</i>	Ni	5,500 mg kg ⁻¹	Robinson <i>et al.</i> (1997)
<i>Eichornia crassipes</i>	Cr (Vi)	6000 mg kg ⁻¹	Lytle <i>et al.</i> (1998)
<i>Iberis intermedia</i>	Ti	3,070 mg kg ⁻¹	Leblanc <i>et al.</i> (1999)
<i>Alternanthera sessilis</i>	Cv	1017 mg kg ⁻¹	Sinha <i>et al.</i> (2002)
<i>Zea mays</i> L. Cv Ganga 5	Cr	2538 mg kg ⁻¹	Sharma <i>et al.</i> (2003)
<i>Pteris vittata</i>	As	23,000 µg g ⁻¹	Dong (2005)
<i>Sesbania drummondii</i>	Cd	1687 mg kg ⁻¹	Israr <i>et al.</i> (2006)
<i>Thlaspi caerulescens</i>	Zn	19410 mg kg ⁻¹	Banasova and Horak (2008)
<i>Thlaspi caerulescens</i>	Cd	80 mg kg ⁻¹	Banasova and Horak (2008)
<i>Myriophyllum heterophyllum</i>	Cd	21.46 µg g ⁻¹	Sivaci <i>et al.</i> (2008)
<i>Potamogeton crispus</i>	Cd	49.09 µg g ⁻¹	Sivaci <i>et al.</i> (2008)
<i>Sorghum sudanense</i>	Cu	5330 mg kg ⁻¹	Wei <i>et al.</i> (2008)
<i>Phragmites australis</i>	Cr	4825 mg kg ⁻¹	Calheiros <i>et al.</i> (2008)
<i>Arabis paniculata</i>	Cd	1127 mg kg ⁻¹ in the shoots	Zeng <i>et al.</i> (2009)
<i>Atriplex halimus</i> subsp. <i>schweinfurthii</i>	Cd	606.51 µg g ⁻¹ DW	Nedjimi and Daoud (2009).
<i>Sedum alfredii</i>	Cd	2,183	Jin and Liu (2009)
<i>Sedum alfredii</i>	Zn	13,799 mg kg ⁻¹ DW	Jin and Liu (2009)
<i>Phytolacca americana</i>	Mn	32,000 µg g ⁻¹	Pollard <i>et al.</i> (2009).
<i>Brassica juncea</i>	Ni	3916 mg kg ⁻¹ dry wt.	Saraswat and Rai (2009)
<i>Potentilla griffithii</i>	Zn	leaves (19,600) mg kg ⁻¹ dry weight	Hu <i>et al.</i> (2009)
<i>Rorippa globosa</i> (Turcz.)	Cd	218.9 µg g ⁻¹ dry weight (DW)	Sun <i>et al.</i> (2010)
<i>Thlaspi praecox</i> Wulfen	Cd	> 1,000 µg g ⁻¹ dry weight in seeds	Vogel-Mikus <i>et al.</i> (2010)

Table 4: Concentrations of heavy metals accumulated in *Vetiveria zizanioides* roots and shoots (Truong, 1999)

Heavy metal	Soil (mg kg ⁻¹)	Shoot (mg kg ⁻¹)	Root (mg kg ⁻¹)	Shoot/root (%)	Shoot/total (%)
As	620.00	11.20	268.0	4.2	4.0
Cd	1.66	0.31	14.2	2.2	2.1
Cu	50.00	13.00	68.0	19.0	16.0
Cr	600.00	18.00	1750.0	1.0	1.0
Pb	730.00	78.20	87.8	87.0	47.0
Hg	6.17	0.12	10.8	11.0	6.0
Ni	300.00	448.00	1040.0	43.0	30.0
Se	74.30	11.30	24.8	46.0	44.0

(Danh *et al.*, 2009). This plant has an ability to accumulate multiple heavy metals in roots and shoots (Table 4) which has been experimentally determined.

The multiple metal accumulation and metal transfer factors from soil (TFS) of three wild macrophyte species viz., *Ipomea* sp., *Eclipta* sp. and *Marsilea* sp. have been studied by Gupta *et al.* (2008) and recorded that *Ipomea* sp. shows transfer factor (TFS) for Cd, Cu, Mn and Zn, while *Eclipta* sp. and *Marsilea* sp. shows TFS ≥1 for Fe, Cu and Cd. The ratio of metals between soil and plant parts (TFS) is an important criterion for the selection of model plant species for phytoremediation and the ratio >1 means higher accumulation of metals in plant parts than soil (Barman *et al.*, 2000). Among aquatic plants, *Elodea densa* showed a high accumulation of mercury in leaves, stems and roots from the natural sediment enriched with CH₃HgCl (Ochiai, 1987) which make this plant potentials for Hg remediation.

PHYSIOLOGY OF METAL HYPERACCUMULATION

Metals influenced the physiology of plants by promote or inhibit the growth. Some metals that required in high concentration suggest a structural or osmotic role, while effects at low concentration may indicate a role as cofactor for specific enzymes. The summary of the beneficial effects of Al, Co, Na, Se and Si on plants under represented (Fig. 1) adopted from Pilon-Smits *et al.* (2009).

Many plants developed Al tolerance characteristics via either apoplastic or symplastic detoxification mechanisms (Ma *et al.*, 2001; Pilon-Smits *et al.*, 2009).

In higher plants, Co has been reported to strongly bind to roots and to be mainly absorbed from the soil solution through passive transport. Co is chemically similar with nickel (Ni) and these two elements enter inside the cells through plasma membrane carriers and may be transported by IRT1 (Pilon-Smits *et al.*, 2009). A strong adsorption of Cd on root apoplast might act as a driving force to extract the metal from the soil, compete with the symplastic absorption and contribute to the amount of metals taken up by the hyperaccumulator, at least in its roots (Redjala *et al.*, 2009).

Se hyperaccumulators namely *Stanleya* sp. and *Astragalus* sp. can accumulate 1000-15000 ppm (0.1-1.5% Se), even from low external concentrations and this has been enhanced by some specialized transporter. Hyperaccumulators such as *Astragalus bisulcatus*, *Brassica oleracea* have a specific selenocysteine methyl transferase, lead to accumulation of Se (Tamaoki *et al.*, 2008)

Toxic metal ions Hg preferentially binds with sulphur and nitrogen rich ligands (amino acids) and entered inside the cells. Hg effect damage include blocking functional groups of enzymes, polynucleotide's, or transport systems for nutrient ions, denaturing and inactivating enzymes and disrupting cell and organelle membrane integrity (Ochiai, 1987). The possible causal mechanisms




Fig. 1: The mechanisms responsible for the growth promoting effects of the five beneficial elements Al, Co, Na, Se and Si (Pilon-Smits *et al.*, 2009)

of Hg toxicity are changes in the permeability of the cell membrane, reactions of sulphhydryl (-SH) groups with cations, affinity for reacting with phosphate groups and active groups of ADP or ATP and replacement of major cations (Kabata-Pendias and Pendias, 1989).

Heavy metals like copper (Cu) and iron (Fe) can be toxic in excess amount because of their participation in redox cycles producing hydroxyl radicals which are extremely toxic to living cells (Stohs and Bagchi, 1995). Unlike Cu and Fe, Cd is a non-redox metal that is strongly phytotoxic and caused growth inhibition and plant death. Cd induced changes in lipid profile (Ouariti *et al.*, 1997) and by affecting the enzymatic activities associated with membranes, such as the H⁺-ATPase (Fodor *et al.*, 1995). Cd is also reported to damage the photosynthetic apparatus (Siedlecka and Baszynsky, 1993), decrease chlorophyll content and inhibit the stomatal regulations (Barcelo and Poschenrieder, 1990). The major storage site for Zn and Cd in plants is cell wall of roots, vacuoles of epidermis and bundle sheath of leaves (Hu *et al.*, 2009). The Cd influx and efflux in leaf mesophyll layer depends on the expression of plasma membrane and tonoplast transporters. Root metal uptake rates are increased, generally through an increase in V_{MAX} , without major changes in K_M , of root metal uptake rates (Lasat *et al.*, 1996; Lombi *et al.*, 2001).

Works on *Alyssum lesbiacum* has given a base of Nickel uptake into vacuoles from leaf tissue and this was enhanced by the presence of Mg/ATP, presumably via energisation of the vacuolar H⁺-ATPase (Ingle and Fricker, 2008). The model hyperaccumulator *Thlaspi caerulescens* is capable to tolerate Zn, Cd and Ni and it has been shown that part of the Ni is translocated as a stable Ni-NA complex in the xylem sap (Mari *et al.*, 2006).

BIOMASS PRODUCTION

The discussion of the effect of heavy metal on biomass productions of metal hyperaccumulation is restricted here to Cd, Pb, Zn, V and Cr.

In metal hyperaccumulators the biomass production level depends on the concentration of the metals and duration of exposures e.g., The biomass is negatively correlate with Cd concentration in *B. napus*, cultured in the nutrient solution and the reduction in dry weight was significantly higher for the root than the shoot. Considering the high aboveground biomass production and Cd accumulation in the shoot, *B. napus* can be a potential candidate for the phytoextraction of Cd (Selvam and Wong, 2008).

Biomass production were extensively studied in split pots filled with soil spiked at 0, 3, 6, 12, 25 and 50 mg Cd kg⁻¹ soil by Selvam and Wong (2009) and revealed decline in biomass production which established that cadmium is toxic for biomass production.

Suitable levels of Pb, Zn and Cd could stimulate the biomass production in *A. paniculata* and thus, it provides a new plant material for understanding the mechanisms of stimulatory effect and co-hyperaccumulation of multiple heavy metal (Tang *et al.*, 2009). However, biomass production is inhibited in *Glycine max* and *Phaseolus vulgaris* in treated with VOSO₄ (Kaplan *et al.*, 1990). Similarly in *E. fluctuans*, biomass production was not affected up to concentration of 2.5 mg L⁻¹ V₂O₅ when exposed for 7 days (Fig. 2). However, V₂O₅ concentration >2.5 mg L⁻¹ significantly reduced the biomass and increased in treatment duration enhanced the V₂O₅ toxicity (Sarma *et al.*, 2009). The author recorded that after 21 days exposure of 10.0 mg L⁻¹ V₂O₅ to *E. fluctuans* approximately 42.47% biomass was decreased (Fig. 2). Similarly in *Vallisneria spiralis* L., an increased in treatment duration enhanced the chromium toxicity and 0.1 µg mL⁻¹ Cr caused 7% decrease in biomass after 48 h and 64% loss of biomass was recorded after 72 h exposure of 10 µg mL⁻¹ Cr (Vajpayee *et al.*, 2001). The results addressed that heavy metals toxicity to biomass




Fig. 2: Biomass production of vanadium treated *E. fluctuans* was negatively correlated with concentration of vanadium in nutrient medium with increased in duration

were concentration and duration's dependent. However, modern transgenic research approach at present going on for introduction of novel traits into high biomass plants for development of effective phytoremediation technologies. A number of transgenic high biomass yield plants have been successfully generated in an attempt to modify the tolerance, uptake or homeostasis of trace elements (Kramer and Chardonnens, 2001).

BIOCHEMICAL RESPONSES

Several heavy metals such as Fe, Cu, Co, Mn, Mo and Ni are essential elements to plant metabolism. In higher concentrations, many heavy metals inhibit plants biochemical production and this has been extensively studied and reviewed (Fernandes and Henriques, 1991; Sarma and Sarma, 2007; Sarma *et al.*, 2009). Photosynthetic pigments of plants belonging to different group's exhibit differential tolerance to metals (Vajpayee *et al.*, 2001). Heavy metal substituted chlorophylls and related porphyrins have been known in vitro for a long time (Kupper *et al.*, 2000). Many researchers examined the effect of heavy metals on photosynthesis and observed a decrease in fluorescence (Atal *et al.*, 1991; El-Sheekh, 1992). Cd induced reduction of photosynthetic pigment were recorded in two species viz., *M. heterophyllum* and *P. crispus* (Fig. 3a, b and 4a, b). The highest decrease in chlorophyll a was recorded in 7.34 mg g⁻¹ in *M. heterophyllum* and 8.09 mg g⁻¹ in *P. crispus* (at 64 mg L⁻¹ and 96 h) have given as evidence for the Cd toxicity to chlorophyll.

Cadmium hyperaccumulator *Atriplex halimus* subsp. *schweinfurthii* was sensitive to high Cd results reduction of chlorophyll pigments, stomatal transpiration and root hydraulic conductivity (Nedjimi and Daoud, 2009). The Cr uptake by many aquatic plants influenced in biochemical process results alteration of pigments and amino acids. It has been reported that Cr (VI) causes toxicity to δ -amionolevulinic acid dehydratase (an enzyme involved in Chlorophyll biosynthesis) by impairing δ -amiono levulinic acid (ALA) utilizations (Vajpayee *et al.*, 2000). Further, Vajpayee *et al.* (2000) suggested that Cr (VI) could exchange the Mg from active site of enzyme resulting into phaeophytin and thus depleted chlorophyll contents in Cr treated plants. Chromium also inhibits chlorophyll biosynthesis by creating nutrient imbalance (Barcelo *et al.*, 1986). It has been reported that chromium induced degradation of carotenoid in some plants while in contrast Vajpayee *et al.* (2001) reported that carotenoid contents was increased in Cr treated *V. spiralis*. Thus, the effects of heavy metals on carotenoid contents were plants and metal specific.




Fig. 3: Chlorophyll a, b and carotenoid contents of (mg g⁻¹ FW) *M. heterophyllum* exposed to various Cd concentrations for (a) 24 and (b) 96 h (Sivaci *et al.*, 2008)




Fig. 4: Chlorophyll a, b and carotenoid contents of (mg g⁻¹ FW) *P. crispus* exposed to various Cd concentrations for (a) 24 and (b) 96 h (Sivaci *et al.*, 2008)

Mercuric cations have a high affinity for sulphhydryl (-SH). In almost all proteins contain sulphhydryl groups or disulphide bridges, Hg could disturb the normal functions of proteins in binding in two sites of a protein molecule without deforming the chain, lead to protein precipitation. (Clarkson, 1972).

Mercury affects both light and dark reactions in photosynthesis and caused inhibitions of electron transport activity, oxygen evolution and quenching of chlorophyll fluorescence in photosystem II (PS II). Substitution of the central atom of chlorophyll, magnesium, by mercury in vivo is an important damage mechanism, because it prevents photosynthetic light harvesting in the affected chlorophyll molecules and results in the breakdown of photosynthesis (Krupa and Baszynski, 1995).

GENETIC AND MOLECULAR BASIS OF METAL HYPERACCUMULATION

Metal hyperaccumulation is a fascinating phenomenon, which has interested scientists for over a century. Hyperaccumulators constitute a group of exceptional plant species and they possess genetically inherited traits of metals hyperaccumulation and tolerance. The understanding of metal hyperaccumulation physiology has recently improved as a result of the development of molecular tools (Verbruggen *et al.*, 2009). Transgenic approaches successfully employed to promote phytoextraction of metals (Cd, Pb and Cu) and metalloids (As, Se) from contaminated soil by their accumulation in the aboveground biomass involved mainly implementation of metal transporters, improved production of enzymes of sulphur metabolism and production of metal-detoxifying chelators metallothioneins and phytochelatin (Kotrba and Najmanova, 2009).

Recent research revealed that *Arabidopsis thaliana* has eight genes encoding members of the type 1 B heavy metal transporting subfamily of the P-type ATPases. Three of these transporters,

HMA2, HMA3 and HMA4, are closely related to each other and are most similar in sequence to the divalent heavy metal cation transporters of prokaryotes (Hussain *et al.*, 2004).

Quantitative mRNA *in situ* hybridization (QISH) in *Thlaspi caerulescens* shows that transporter gene expression changes during cadmium (Cd)/zinc (Zn) hyperaccumulations. Members of the ZIP gene family, a novel metal transporter family first time identified in plants are capable of transporting a variety of cations including Cd, Fe, Mn and Zn. The different cellular expression patterns for ZNT1 and ZNT5 were recorded by Kupper and Kochian (2010) both belonging to the ZIP family of transition metal transporters. ZNT1 may function in micronutrient nutrition while ZNT5 may be involved in metal storage associated with hyperaccumulation. Cadmium induced changes in cellular expression for ZNT1, ZNT5 and MTP1 could also be part of plants acclimatization to Cd toxicity.

The integral membrane protein *Thlaspi goesingense* metal tolerance protein 1 (TgMTP1) has been suggested to play an important role in Zn hyperaccumulation. High levels of TgMTP1 at the vacuolar membrane in shoot tissue of the Zn hyperaccumulator *T. goesingense* has been played an important role in both Zn tolerance and enhanced Zn uptake and accumulation, via the activation of a systemic Zn deficiency response (Gustin *et al.*, 2009).

An antiporter is an integral membrane protein which is involved in secondary active transport of two or more ions across a phospholipid membrane in opposite directions. A recent genomic analysis provides a breakthrough in acquisition of zinc hypertolerance and hyperaccumulation characters via involvement of Zn²⁺/H⁺ antiporter. Metal Tolerance Protein 1 (MTP1) is one of genes present in *A. halleri* which encodes a Zn²⁺/H⁺ antiporter involved in cytoplasmic zinc detoxification and developed a zinc tolerance characteristic (Shahzad *et al.*, 2010).

Plant tissue cultures such as callus, cell suspensions and hairy roots are extensively used in phytoremediation research as model plant systems. Many studies have demonstrated that plant tissue cultures are an extremely valuable tool in phytoremediation research. The results derived from tissue cultures can be used to predict the responses of plants to environmental contaminants and to improve the design and thus reduce the cost of subsequent conventional whole plant experiments (Doran, 2009).

ANTIOXIDANTS DEFENSE SYSTEM IN METAL HYPERTOLERANCE

Phytochelatins are oligomers of glutathione, produced by the enzyme phytochelatase. They are found in plants, fungi, nematodes and all groups of algae including cyanobacteria. Phytochelatins act as chelators and are important for heavy metal detoxification (Ha *et al.*, 1999). They are abbreviated PC2 through PC11. Plants possess several antioxidative defense systems to scavenge toxic free radicals in order to protect themselves from the oxidant stress including that caused by heavy metals. The antioxidative defense system falls into two general classes: (1) low molecular weight antioxidants, which consist of lipid-soluble membrane-associated antioxidants (e.g., α -tocopherol and β -carotene) and water-soluble reductants (e.g., glutathione and ascorbate) and (2) antioxidative enzymes: Superoxide Dismutase (SOD) Ascorbate Peroxidase (APX), Catalase (CAT) and Glutathione Reductase (GR).

Glutathione (GSH), a sulfur containing tripeptide, is considered to be the most important cellular antioxidant involved in cellular defense against toxicants (Scott *et al.*, 1993) and function directly as a free radical scavenger. This antioxidant is also the precursor for the phytochelatins that act as heavy metal binding peptides in plants (Rosen, 2002). GSH levels in plants are known to change under metal stress (Koricheva *et al.*, 1997).

The Cd treatments significantly enhanced the synthesis of phytochelatins (PCS) in plants. However, Sun *et al.* (2010) reported that the variation in phytochelatins productions in root and shoots in two Cd treated species viz., *R. globosa* and *R. islandica* may be used as a biomarker of Cd hyperaccumulation and the synthesis of PCS may be related to an increase in the uptake of Cd ions into the cytoplasm, not the primary mechanism for Cd tolerance. Similarly the accumulation of Cd has influenced on the synthesis of phytochelatins in *Brassica napus* and in the shoot, the concentration of PC3 and PC4 was higher than the PC2 irrespective of the quantity of Cd uptake (Selvam and Wong, 2008). This result gives an evidences that the detoxification of Cd involves higher molecular weight thiol complexes in the shoot. In *Arabis paniculata* after Cd exposure induced formation of PCS and three unknown thiols in the roots, but none were detected in the shoots (Zeng *et al.*, 2009). *Sesbania* sp. responded to Hg induced oxidative stress by modulating non-enzymatic antioxidants viz., Glutathione (GSH) and Non-Protein Thiols (NPSH) and enzymatic antioxidants: Superoxide Dismutase (SOD), ascorbate peroxidase (APX) and Glutathione Reductase (GR). The *Sesbania* plants were able to tolerate Hg induced stress using effective antioxidative defense mechanisms (Israr *et al.*, 2006; Israr and Sahi, 2006). It has been suggested that phytochelatins play a constitutive role in plant metal tolerance (Zenk, 1996). However, evidence provided by Zenk (1996) for this role is not conclusive and could just as easily indicate a stress response (e.g., production of phytochelatins upon exposure to metals).

CONCLUSION

Phytoremediation is initiated all over the globe and this has considered one of the low-cost novel green technologies. The physiological and molecular basis of metal hyperaccumulation in plants are still in research and development phase. This review has focused on recent evidence that identifies potential molecular mechanism that may be involved in the resistance, tolerance as well as hyperaccumulation of heavy metals. The findings suggest that in some plants ZIP family genes contribute to metal hyperaccumulation and transport, but their individual functions yet to be identified and further intensive research is needed in this concern. The identifications of individual functions of metal transporters will help to develop the knowledge of plants metal homeostasis. Results already obtained have been indicated that the physiological and biochemical responses were plants and metal specific. Identification of novel genes with high biomass yield characteristics and the subsequent development of transgenic plants with superior remediation capacities will be encouraging further research. In depth research study is warranted to find out which plant is maximum resistant and best adapted in particular metallic environment or region. In situ toxicity test could be beneficial for initial identification of particular species.

ACKNOWLEDGMENTS

I am thankful to all colleagues, in particular to current and previous members of my research team, for their supports. I thank to all my Lab mates of Ecosystem Research Laboratory, North-Eastern Hill University. I am grateful to Ms. Idaiarilin Lyndogh Synrem for reading the first draft of the manuscripts and Mr. Souravjyoti Borah for technical assistant. Funding by a Dr. D.S. Kothari Post Doctoral Fellowship from the University Grants Commission, New Delhi (F. 4-2/2006 (BSR)/13-177/2008 (BSR) is gratefully acknowledged. Finally, I apologize to all colleagues for those parts of their work that were not cited in the manuscripts because of restrictions in space.

REFERENCES

- Adriano, D.C., 1986. Trace Elements in the Terrestrial Environment. 1st Edn., Springer-Verlag, New York.
- Assuncao, A.G.L and H. Schat, 2003. *Thlaspi caerulescens*, an attractive model species to study heavy metal hyperaccumulation in plants. *New Phytol.*, 159: 351-360.
- Atal, N., P.P. Saradhi and P. Mohanty, 1991. Inhibition of the chloroplast photochemical reactions by treatment of wheat seedlings with low concentrations of cadmium: analysis of electron transport activities and changes in fluorescence yield. *Plant Cell Physiol.*, 32: 943-951.
- Baker, A.J.M, R.D. Reeves and A.S.M. Hajar, 1994. Heavy metal hyperaccumulation and tolerance in British populations of the metallophyte *Thlaspi caerulescens* J. and C. Presl (Brassicaceae). *New Phytol.*, 127: 61-68.
- Baker, A.J.M. and P.L. Walker, 1989. Ecophysiology of Metal Uptake by Tolerant Plants. In: Heavy Metal Tolerance in Plants: Evolutionary Aspects, Shaw, A.J. (Ed.), CRC. Publ., Boca Raton, FL, pp: 155-177.
- Baker, A.J.M. and R.R. Brooks, 1989. Terrestrial higher plants which hyper accumulate metallic elements -Review of their distribution, ecology and phytochemistry. *Biorecovery*, 1: 81-126.
- Baker, A.J.M., R.D. Reeves and S.P. McGrath, 1991. *In situ* Decontamination of Heavy Metal Polluted Soils Using Crops of Metal Accumulating Plants -A Feasibility Study. In: *In situ* Bioremediation, Hinchee, R.E. and R.F. Olfenbuttel (Eds.), Butterworth-Heinemann, Stoneham, M.A., pp: 539-544.
- Baker, A.J.M., S.P. McGrath, R.D. Reeves and J.A.C. Smith, 2000. Metal Hyperaccumulator Plants: A Review of the Ecology and Physiology of a Biological Resource for Phytoremediation of Metal Polluted Soils. In: Phytoremediation of Contaminated Soil and Water, Terry, N. And G. S. Banuelos (Eds.), CRC Press, Boca Raton, pp: 85-107.
- Banasova, V. and O. Horak, 2008. Heavy metal content in *Thlaspi caerulescens* J. et C. Presl growing on metalliferous and non-metalliferous soils in Central Slovakia. *Int. J. Environ. Pollut.*, 33: 133-145.
- Barcelo, J. and C. Poschenrieder, 1990. Plant water relations as affected by heavy metal stress: A review. *J. Plant Nutr.*, 13: 1-37.
- Barcelo, J., C. Poschenrieder and B. Gunse, 1986. Water relations of chromium VI treated bush bean plants (*Phaseolus vulgaris* L. cv. Contender) under both normal and water stress conditions. *J. Exp. Bot.*, 37: 178-187.
- Barman, S.C., R.K. Sahu, S.K. Bhargava and C. Chatterjee, 2000. Distribution of heavy metals in wheat, mustard and weed grown in fields irrigated with industrial effluents. *Bull. Environ. Contam. Toxicol.*, 64: 489-496.
- Barman, S.C., G.C. Kisku, P.R. Salve, D. Misra, R.K Sahu, P.W. Ramteke and S.K. Bhargava, 2001. Assessment of industrial effluent and its impact on soil and plants. *J. Environ. Biol.*, 22: 251-256.
- Baudouin, C., M. Charveron, R. Tarrouse and Y. Gall, 2002. Environmental pollutants and skin cancer. *Cell. Biol. Toxicol.*, 18: 341-348.
- Beath, O.A., H.F. Eppson and G.S. Gilbert, 1937. Selenium distribution in and seasonal variation of vegetation type occurring on seleniferous soils. *J. Am. Pharma. Assoc.*, 26: 394-405.
- Bert, V. and P. Meerts, 2003. Genetic basis of Cd tolerance and hyperaccumulation in *Arabidopsis halleri*. *Plant Soil*, 249: 9-18.

- Beyersmann, D and A. Hartwig, 2008. Carcinogenic metal compounds: Recent insight into molecular and cellular mechanisms. *Arch. Toxicol.*, 82: 493-512.
- Blaylock, M.J., D.E. Salt, S. Dushenkov, O. Zakhrova, C. Gussman and Y. Kapulnik, 1997. Enhanced accumulation of Pb in Indian mustard by soil-applied chelating agents. *Environ. Sci. Technol.*, 31: 860-865.
- Boonyapookana, B., P. Parkplan, S. Techapinyawat, R.D. DeLaune and A. Jugsujinda, 2005. Phytoaccumulation of lead by sunflower (*Helianthus annuus*), tobacco (*Nicotiana tabacum*) and vetiver (*Vetiveria zizanioides*). *J. Environ. Sci. Health A Tox. Hazard. Subst. Environ. Eng.*, 40: 117-137.
- Calheiros, C.S.C., A.O.S.S. Rangel and P.M.L. Castro, 2008. The effects of tannery wastewater on the development of different plant species and chromium accumulation in phragmites australis. *Arch. Environ. Contam. Toxicol.*, 55: 404-414.
- Cardwell, A.J., D.W. Hawker and M. Greenway, 2002. Metal accumulation in aquatic macrophytes from southeast Queensland, Australia. *Chemosphere*, 48: 653-663.
- Chandra Sekhar, K., C.T. Kamala, N.S. Chary, V. Balaram and G. Garcia, 2005. Potential of *Hemidesmus indicus* for phytoextraction of lead from industrially contaminated soils. *Chemosphere*, 58: 507-514.
- Clarkson, T.W., 1972. The pharmacology of mercury compounds. *Ann. Rev. Pharmacol.*, 12: 375-406.
- Dan, T.V., S. Krishna Raj and P.K. Saxena, 2000. Metal tolerance of scented geranium (*Pelargonium* sp. Frensham): Effects of cadmium and nickel on chlorophyll fluorescence kinetics. *Int. J. Phytorem.*, 2: 91-104.
- Dan, T.V., S. Krishnaraj and P.K. Saxena, 2002. Cadmium and nickel uptake and accumulation in scented geranium (*Pelargonium* sp. frensham). *Water Air Soil Pollut.*, 137: 355-364.
- Danh, L.T., P. Truong, R. Mammucari, T. Tran and N. Foster, 2009. Vetiver grass, *Vetiveria zizanioides*: A choice plant for phytoremediation of heavy metals and organic wastes. *Int. J. Phytorem.*, 11: 664-691.
- Davis, M.A. and S.G. Pritchard, R.S Boyd and S.A. Prior, 2001. Developmental and induced responses of nickel-based and organic defences of the nickel-hyperaccumulating shrub, *Psychotria douarrei*. *New Phytol.*, 150: 49-58.
- Deng, D.M., W.S. Shu, J. Zhang, H.L. Zou, Z.H. Ye, M.H. Wong and Z. Lin, 2007. Zinc and cadmium accumulation and tolerance in populations of *Sedum alfredii*. *Environ. Pollut.*, 147: 381-386.
- Dong, R., 2005. Molecular cloning and characterization of a phytochelatin synthase gene, PvPCS1, from *Pteris vittata* L. *J. Ind. Microbiol. Biot.*, 32: 527-533.
- Doran, P.M., 2009. Application of plant tissue cultures in phytoremediation research: Incentives and limitations. *Biotechnol. Bioeng.*, 103: 60-76.
- Ebbs, S.D. and L.V. Kochian, 1997. Toxicity of Zn and Copper to *Brassica* species: Implication for phytoremediation. *J. Environ. Qual.*, 26: 776-781.
- El-Sheekh, M.M., 1992. Inhibition of photosystem II in the green alga *Scenedesmus obliquus* by nickel. *Biochem. Physiol. Pfl.*, 188: 363-372.
- Ellis, D.R., T.G. Sors, D.G. Brunk, C. Albrecht and C. Orser *et al.*, 2004. Production of Se-methylselenocysteine in transgenic plants expressing selenocysteine methyltransferase. *BMC Plant. Biol.*, 4: 1-15.

- Falandysz, J., K. Lipka, M. Kawano, A. Brzostowski, M. Dadey, A. Jedrusiak and T. Puzyn, 2003. Mercury content and its bio-concentration factors in wild mushrooms at lukta and morag, north-eastern Poland. *J. Agric. Food. Chem.*, 51: 2832-2836.
- Fernandes, J.C. and F.S. Henriques, 1991. Biochemical, physiological and structural effects of excess copper in plants. *Bot. Rev.*, 57: 246-273.
- Flathman, P.E. and G.R. Lanza, 1998. Phytoremediation: Current views on an emerging green technology. *Soil Sediment Contamin. Int. J.*, 71: 415-432.
- Fodor, E., A. Szabo-Nagy and L. Erdei, 1995. The effects of cadmium on the fluidity and H⁺-ATPase activity of plasma membrane from sunflower and wheat roots. *J. Plant. Physiol.*, 147: 87-92.
- Garcia, G., A. Faz and M. Cunha, 2004. Performance of *Piptatherum miliaceum* (Smilo grass) in edaphic Pb and Zn phytoremediation over a short growth periods. *Int. Biodeters. Biodeg.*, 54: 245-250.
- Glass, D.J., 1999. U.S. and International Markets for Phytoremediation, 1999-2000. D. Glass Associates, Needham MA.
- Gleba, D., N.V. Borisjuk, L. Borisjuk, R. Kneer and A. Poulev *et al.*, 1999. Use of plant roots for phytoremediation and molecular farming. *Proc. Natl. Acad. Sci.*, 96: 5973-5977.
- Gupta, S., S. Nayek, R.N. Saha and S. Satpati, 2008. Assessment of heavy metal accumulation in macrophyte, agricultural soil and crop plants adjacent to discharge zone of sponge iron factory. *Environ. Geol.*, 55: 731-739.
- Gustin, J.L., M.E. Loureiro, D. Kim, G. Na, M. Tikhonova and D.E. Salt, 2009. MTP1-dependent Zn sequestration into shoot vacuoles suggests dual roles in Zn tolerance and accumulation in Zn-hyperaccumulating plants. *Plant J.*, 57: 1116-1127.
- Ha, S.B., A.P. Smith, R. Howden, W.M. Dietrich and S. Bugg *et al.*, 1999. Phytochelatin synthase genes from arabidopsis and the yeast *Schizosaccharomyces pombe*. *Plant Cell*, 11: 1153-1164.
- Hu, P.J., R.L. Qiu, P. Senthilkumar, D. Jiang, Z.W. Chen, Y.T. Tang and F.J. Liu, 2009. Tolerance, accumulation and distribution of zinc and cadmium in hyperaccumulator *Potentilla griffithii*. *Environ. Exp. Bot.*, 66: 317-325.
- Huang, J.W., J. Chen, W.R. Berti and S.D. Cunningham, 1997. Phytoremediation of lead contaminated soil: Role of synthetic chelates in lead phytoextraction. *Environ. Sci. Technol.*, 31: 800-805.
- Hussain, D., M.J. Haydon, Y. Wang, E. Wong and S.M. Sherson *et al.*, 2004. P-Type ATPase Heavy Metal Transporters with Roles in Essential Zinc Homeostasis in Arabidopsis. *Plant Cell*, 16: 1327-1339.
- Ingle, R.A. and M.D. Fricker, 2008. Evidence for nickel/proton antiport activity at the tonoplast of the hyperaccumulator plant *Alyssum lesbiacum*. *Plant Biol.*, 10: 746-753.
- Israr M., S.V. Sahi and J. Jain, 2006. Cadmium accumulation and antioxidative responses in the sesbania *Drummondii callus*. *Arch. Environ. Contam. Toxicol.*, 50: 121-127.
- Israr, M. and S. Sahi, 2006. Bioaccumulation and physiological effects of mercury in sesbania drummondii. *Chemosphere*, 65: 591-598.
- Jaffre, T., R.R. Brooks, J. Lee and R.D. Reeves, 1976. *Sebertia acumip* A nickel-accumulating plant from New Caledonia. *Science*, 193: 579-580.
- Jain, S.K., P. Vasudevan and N.K. Jha, 1990. *Azolla pinnata* r.br. and *Lemna minor* l. for removal of led and Zn from polluted water. *Water Res.*, 24: 177-183.

- Jin, X.F. and D. Liu, 2009. Effects of zinc on root morphology and antioxidant adaptations of cadmium-treated *Sedum alfredii* H. *J. Plant Nutr.*, 32: 1642-1656.
- Kabata-Pendias, A. and H. Pendias, 1989. Trace Elements in Soil and Plants. CRC Press, Boca Raton, FL.
- Kaplan, D.I., D.C. Adriano, C.L. Carlson and S. Sajwan, 1990. Vanadium: Toxicity and accumulation by Beans. *Water Air Soil Pollut.*, 49: 81-91.
- Karimi, N., S.M. Ghaderian, A. Raab, J. Feldmann and A.A. Meharg, 2009. An arsenic-accumulating, hypertolerant *Brassica isatis*, *Cappadocica*. *New Phytol.*, 184: 41-47.
- Kim, S.G., J.H. Jee and J.C. Kang, 2004. Cadmium accumulation and elimination in tissues of juvenile olive flounder, *Paralichthys olivaceus* after sub-chronic cadmium exposure. *Environ. Pollut.*, 127: 117-123.
- Knasmuller, S., E. Gottmann, H. Steinkellner, A. Fomin, C. Pickl, A. Paschki, R. God and M. Kundi, 1998. Detection of genotoxic effects of heavy metal contaminated soils with plant bioassays. *Mutat. Res.*, 420: 37-48.
- Koricheva, J., S. Roy, J.A. Vranjic, E. Haukioja, P.R. Hughes and O. Hanninen, 1997. Antioxidants responses to stimulated acid rain and heavy metal deposition in birch seedlings. *Environ. Pollut.*, 95: 249-258.
- Kotrba, P. and J. Najmanova, 2009. Genetically modified plants in phytoremediation of heavy metal and metalloid soil and sediment pollution. *Biotechnol. Adv.*, 27: 799-810.
- Kramer, U., J.D. Cotter-Howells, J.M. Charnock, A.J.M. Baker and J.A.C. Smith, 1996. Free histidine as a metal chelator in plants that accumulate nickel. *Nature*, 379: 635-638.
- Kramer, U. and A.N. Chardonens, 2001. The use of transgenic plants in the bioremediation of soils contaminated with trace elements. *Applied Microbiol. Biotechnol.*, 55: 661-672.
- Kramer, U., 2010. Metal hyperaccumulation in plants. *Annu. Rev. Plant Biol.*, 61: 517-534.
- KrishnaRaj, S., T. Dan and P. Saxena, 2000. A fragrant solution to soil remediation. *Int. J. Phytorem.*, 2: 117-132.
- Krupa, Z. and T. Baszynski, 1995. Some aspects of heavy metals toxicity towards photosynthetic apparatus: Direct and indirect effects on light and dark reactions. *Acta. Physiol. Plant.*, 17: 177-190.
- Kubota, H. and C. Takenaka, 2003. *Arabis gemmifera* is a hyperaccumulator of Cd and Zn. *Int. J. Phytorem.*, 5: 197-220.
- Kulakow, P.A., A.P. Schwab and M.K. Banks, 2000. Screening plant species for growth on weathered, petroleum hydrocarbon-contaminated sediments. *Int. J. Phytorem.*, 2: 297-317.
- Kupper, H., E. Lombi, F.J. Zhao and S.P. McGrath, 2000. Cellular compartmentation of cadmium and zinc in relation to other elements in the hyperaccumulator *Arabidopsis halleri*. *Planta*, 212: 75-84.
- Kupper, H. and L.V. Kochian, 2010. Transcriptional regulation of metal transport genes and mineral nutrition during acclimatization to cadmium and zinc in the Cd/Zn hyperaccumulator, *Thlaspi caerulescens* (Ganges population). *New Phytol.*, 185: 114-129.
- Lasat, M.M., A.J.M. Baker and L.V. Kochian, 1996. Physiological characterization of root Zn²⁺ absorption and translocation to shoots in Zn hyperaccumulator and non-accumulator species of *Thlaspi*. *Plant Physiol.*, 112: 1715-1722.
- Lasat, M.M., 2002. Phytoextraction of toxic metals -A review of biological mechanisms. *J. Environ. Qual.*, 31: 109-120.

- Leblanc, M., D. Petit, A. Deram, B. Robinson and R.R. Brooks, 1999. The hyperaccumulation and environmental significance of hyperaccumulation of thallium by *Iberis intermedia* from southern France. *Econ. Geol.*, 94: 109-113.
- Lombi, E., F.J. Zhao, S.P. McGrath, S.D. Young and G.A. Sacchi, 2001. Physiological evidence for a high-affinity cadmium transporter highly expressed in a *Thlaspi caerulescens* ecotype. *New Phytol.*, 149: 53-60.
- Lytle, C.M., F.W. Lytle, N. Yang, Q. JinHong, D. Hansen, A. Zayed and N. Terry, 1998. Reduction of Cr (VI) to Cr (III) by wetland plants: Potential for *in situ* heavy metal detoxification. *Environ. Sci. Technol.*, 32: 3087-3093.
- Ma, L.Q., K.M. Komar, C. Tu, W. Zhang, Y. Cai and E.D. Kennelley, 2001. A fern that hyperaccumulates arsenic. *Nature*, 409: 579-579.
- Manousaki, E., J. Kadukova, N. Papadantonakis and N. Kalogerakis, 2008. Phytoextraction and phytoexcretion of Cd by the leaves of *Tamarix smyrnensis* growing on contaminated non-saline and saline soils. *Environ. Res.*, 106: 326-332.
- Mari, S., D. Gendre, K. Pianelli, L. Ouerdane and R. Lobinski *et al.*, 2006. Root-to-shoot long-distance circulation of nicotianamine and nicotianamine-nickel chelates in the metal hyperaccumulator *Thlaspi caerulescens*. *J. Exp. Bot.*, 57: 4111-4122.
- Mijovilovich, A., B. Leitenmaier, W. Meyer-Klaucke, P.M. Kroneck, B. Götz and H. Kupper, 2009. Complexation and toxicity of copper in higher plants. II. Different mechanisms for copper versus cadmium detoxification in the copper-sensitive cadmium/zinc hyperaccumulator *Thlaspi caerulescens* (Ganges Ecotype). *Plant Physiol.*, 151: 715-731.
- Mizuno, T., K. Hirano, S. Kato and H. Obata, 2008. Cloning of ZIP family metal transporter genes from the manganese hyperaccumulator plant *Chengiopanax sciadophylloides* and its metal transport and resistance abilities in yeast. *Soil Sci. Plant Nutr.*, 54: 86-94.
- Mkandavire, M. and E.G. Dude, 2005. Accumulation of arsenic in *Lemna gibba* L. (duckweed) in tailing waters of two abandoned uranium mining sites in Saxony, Germany. *Sci. Total Environ.*, 336: 81-89.
- Nedjimi, B. and Y. Daoud, 2009. Cadmium accumulation in *Atriplex halimus* subsp. *schweinfurthii* and its influence on growth, proline, root hydraulic conductivity and nutrient uptake. *Flora Morphol. Distribution Funct. Ecol. Plants*, 204: 316-324.
- Ochiai, E.I., 1987. *General Principles of Biochemistry of the Elements*. Plenum Press, New York.
- Odjegba, V.J. and I.O. Fasidi, 2004. Accumulation of trace elements by *Pistia stratiotes*: Implications for phytoremediation. *Ecotoxicology*, 13: 637-646.
- Otte, M.L., M.S. Haarsma, R.A. Broekman and J. Rozema, 1993. Relation between heavy metal concentrations in salt marsh plants and soil. *Environ. Pollut.*, 82: 13-22.
- Ouariti, O., N. Boussama, M. Zarrouk, A. Cherif and M.H. Ghorbali, 1997. Cadmium and copper induced changes in tomato membranes lipids. *Phytochemistry*, 45: 1343-1350.
- Padmavathiamma, P.K. and Y.L. Loretta, 2007. Phytoremediation technology: Hyper-accumulation metals in plants. *Water Air Soil Pollut.*, 184: 105-126.
- Pilon-Smits, E.A.H., C.F. Quinn, W. Tapken, M. Malagoni and M. Schiavon, 2009. Physiological functions of beneficial elements. *Curr. Opin. Plant Biol.*, 12: 267-274.
- Pollard, A.J., H.L. Stewart and C.B. Roberson, 2009. Manganese hyperaccumulation in *Phytolacca americana* L. from the southeastern United States. *Northeastern Natur.*, 16: 155-162.
- Prasad, M.N.V., 2003. Phytoremediation of metal-polluted ecosystems: Hope for commercialization. *Russ. J. Plant Physiol.*, 50: 764-780.

- Prasad, M.N.V. and H. Freitas, 2003. Metal hyperaccumulation in plants: Biodiversity prospecting for phytoremediation technology. *Electron. J. Biotechnol.*, 6: 275-321.
- Rai, U.N., S. Sinha, R.D. Tripathi and P. Chandra, 1995. Waste water treatability potential of some aquatic macrophytes: Removal of heavy metals. *Ecol. Eng.*, 5: 5-12.
- Redjala, T., T. Sterckeman and J.L. Morel, 2009. Cadmium uptake by roots: Contribution of apoplast and of high- and low-affinity membrane transport systems. *Environ. Exp. Bot.*, 67: 235-242.
- Reeves, R.D. and A.J.M. Baker, 2000. Metal-Accumulating Plants. In: *Phytoremediation of Toxic Metals-Using Plants to Clean Up the Environment*, Raskin, I. and B.D. Ensley (Eds.). Wiley, New York, pp: 193-229.
- Reeves, R.D. and R.R. Brooks, 1983. Hyperaccumulation of lead and zinc by two metallophytes from a mining area of central Europe. *Environ. Pollut. A Ecol. Biol.*, 31: 277-287.
- Robinson, B.H., R.R. Brooks, A.W. Howes, J.H. Kirkman and P.E.H. Gregg, 1997. The potential of the high biomass nickel hyperaccumulator *Berkheya coddii* for phytoremediation and phytomining. *J. Geochem. Explor.*, 60: 115-126.
- Rosen, B.P., 2002. Biochemistry of arsenic detoxification. *FEBS Lett.*, 529: 86-92.
- Sahi, S.V., N.L. Bryant, N.C. Sharma and S.R. Singh, 2002. Characterization of a lead hyperaccumulator shrub, *Sesbania drummondii*. *Environ. Sci. Technol.*, 36: 4676-4680.
- Saraswat, S. and J.P.N. Rai, 2009. Phytoextraction potential of six plant species grown in multimetal contaminated soil. *Chem. Ecol.*, 25: 1-11.
- Sarma, H. and C.M. Sarma, 2007. Impact of fertilizer Industry effluent on plant chlorophyll, protein and total sugar. *Nat. Env. Pollut. Technol.*, 6: 633-636.
- Sarma, H., 2008. Effect of air and water pollutants released from Namrup fertilizer complex on vegetation of surrounding area. Ph.D. Thesis, Gauhati University, India.
- Sarma, H., A. Sarma and C.M. Sarma, 2009. Physiological studies of some weeds grown under heavy metal and industrial effluent discharge zone of fertilizer factory. *J. Ecol. Nat. Environ.*, 1: 173-177.
- Scott, N., K.M. Hatlelid, N.E. MacKenzie and D.E. Carter, 1993. Reactions of arsenic (III) and arsenic (V) species with glutathione. *Chem. Res. Toxicol.*, 6: 102-106.
- Selvam, A. and J.W. Wong, 2008. Phytochelatin synthesis and cadmium uptake of *Brassica napus*. *Environ. Technol.*, 29: 765-773.
- Selvam, A. and J.W.C. Wong, 2009. Cadmium uptake potential of *Brassica napus* cocropped with *Brassica parachinensis* and *Zea mays*. *J. Hazard. Mater.*, 167: 170-178.
- Shahzad, Z., F. Gosti, H. Frérot, E. Lacombe, N. Roosens, P. Saumitou-Laprade and P. Berthomieu, 2010. The five AhMTP1 zinc transporters undergo different evolutionary fates towards adaptive evolution to zinc tolerance in *Arabidopsis halleri*. *PloS Genet.*, 6: e1000911-e1000911.
- Sharma, D.M., C.P. Sharma and R.D. Tripathi, 2003. Phytotoxic lesions of chromium in maize. *Chemosphere*, 51: 63-68.
- Sharma, N.C., J.L. Gardea-Torresdey, J. Parsons and S.V. Sahi, 2004. Chemical speciation and cellular deposition of lead in *Sesbania drummondii*. *Environ. Toxicol. Chem.*, 23: 2068-2073.
- Siedlecka, A. and T. Baszynski, 1993. Inhibition of electron flow around photosystem I in chloroplasts of Cd-treated maize plants is due to Cd-induced iron deficiency. *Physiol. Plant*, 87: 199-202.
- Singh, O.V., S. Labana, G. Pandey, R. Budhiraja and R.K. Jain, 2003. Phytoremediation: An overview of metal ion decontamination from soil. *Applied Microbiol. Biotech.*, 61: 405-412.

- Sinha, S., R. Saxena and S. Singh, 2002. Comparative studies on accumulation of Cr from metal solution and tannery effluent under repeated metal exposure by aquatic plants: Its toxic effects. *Environ. Monit. Assess.*, 80: 17-31.
- Sivaci, A., E. Elmas, F. Gumu and E.R. Sivaci, 2008. Removal of cadmium by *Myriophyllum heterophyllum michx* and *Potamogeton crispus* L. and its effect on pigments and total phenolic compounds. *Arch. Environ. Contam. Toxicol.*, 54: 612-618.
- Spinoza-Quinones, F.R., C.E. Zacarkim, S.M. Palacio, C.L. Obregon and D.C. Zenatti *et al.*, 2005. Removal of heavy metal from polluted river water using aquatic macrophytes *Salvinia* sp. *Braz. J. Physiol.*, 35: 744-746.
- Stohs, S.J. and D. Bagchi, 1995. Oxidative mechanism in the toxicity of metal ions. *Free Radic. Biol. Med.*, 18: 321-336.
- Sun, Q., Z.H. Ye, X.R. Wang and M.H. Wong, 2005. Increase of glutathione in mine population of sedum alfredii: A Zn hyperaccumulator and Pb accumulator. *Phytochemistry*, 66: 2549-2556.
- Sun, R., C. Jin and Q. Zhou, 2010. Characteristics of cadmium accumulation and tolerance in *Rorippa globosa* (Turcz.) Thell., a species with some characteristics of cadmium hyperaccumulation. *Plant Growth Regul.*, 61: 67-74.
- Suresh, B. and G.A. Ravishankar, 2004. Phytoremediation-a novel and promising approach for environmental clean-up. *Crit. Rev. Biotechnol.*, 24: 97-124.
- Tamaoki, M., J.L. Freeman and E.A.H. Pilon-Smits, 2008. Cooperative ethylene and jasmonic acid signaling regulates selenite resistance in arabidopsis. *Plant Physiol.*, 146: 1219-1230.
- Tang, Y.T., R.L. Qiu, X.W. Zeng, R.R. Ying, F.M. Yu and X.Y. Zhou, 2009. Lead, zinc, cadmium hyperaccumulation and growth stimulation in *Arabis paniculata* Franch. *Environ. Exp. Bot.*, 66: 126-134.
- Tian, J.L., H.T. Zhu, Y.A. Yang and Y.K. He, 2004. Organic mercury tolerance, absorption and transformation in Spartina plants. *J. Plant Physiol. Mol. Biol.*, 30: 577-582.
- Truong, P., 1999. Vetiver grass technology for mine rehabilitation. *Pacific Rim Vetiver Network Tech. Bull.*, 2: 1-19.
- Truong, P., 2003. Vetiver system for water quality improvement. *Proceeding of 3rd International Vetiver Conference*, Oct. 6-9, Guangzhou, China, pp: 61-74.
- USEPA, 2000. *Introduction to Phytoremediation*. EPA., Washington, DC.
- Vajpayee, P., R.D. Tripathi, U.N. Rai M.B. Ali and S.N. Sinha, 2000. Chromium (VI) accumulation reduced chlorophyll biosynthesis, nitrate reductase activity and protein content of *Nymphaea alba* L. *Chemosphere*, 41: 1075-1082.
- Vajpayee, P., U.N. Rai, M.B. Ali, R.D. Tripathi, V. Yadav and S.N. Sinha, 2001. Chromium induced physiological changes in *Vallisneria spiralis* L. and its role in phytoremediation of tannery effluent. *Bull. Environ. Contam. Toxicol.*, 67: 246-256.
- Verbruggen, N., C. Hermans and H. Schat, 2009. Molecular mechanisms of metal hyperaccumulation in plants. *New Phytol.*, 181: 759-776.
- Vogel-Mikus, K., I. Arcon and A. Kodre, 2010. Complexation of cadmium in seeds and vegetative tissues of the cadmium hyperaccumulator *Thlaspi praecox* as studied by X-ray absorption spectroscopy. *Plant Soil*, 331: 439-451.
- Watanabe, T, M.R. Broadley, S. Jansen, P.J. White and J. Takada *et al.*, 2007. Evolutionary control of leaf element composition in plants. *New Phytol.*, 174: 516-523.
- Wei, S.H., Q.X. Zhou, X. Wang, W. Cao, L.P. Ren and Y.F. Song, 2004. Potential of weed species applied to remediation of soils contaminated with heavy metals. *J. Environ. Sci.*, 16: 868-873.

- Wei, L., C. Luo, X. Li and Z. Shen, 2008. Copper accumulation and tolerance in *Chrysanthemum coronarium* L. and *Sorghum sudanense* L. *Arch. Environ. Contam. Toxicol.*, 55: 238-246.
- Wu, H.B. and S.R. Tang, 2009. Using elevated CO₂ to increase the biomass of a *Sorghum vulgare* x *Sorghum vulgare* var. *sudanense* hybrid and *Trifolium pratense* L. and to trigger hyperaccumulation of cesium. *J. Hazard. Mater.*, 170: 861-870.
- Xian, X. and G. Shokohifard, 1989. Effect of pH on chemical forms and plant availability of cadmium, zinc and lead in polluted soils. *Water Air Soil Pollut.*, 45: 265-273.
- Xiong, Y.H., X.E. Yang, Z.Q. Ye and Z.L. He, 2004. Characteristics of cadmium uptake and accumulation by two contrasting ecotypes of *Sedum alfredii* Hance. *J. Environ. Sci. Health A Tox. Hazard Subst. Environ. Eng.*, 39: 2925-2940.
- Zeng, X., L.Q. Ma, R. Qiu and Y. Tang, 2009. Responses of non-protein thiols to Cd exposure in Cd hyperaccumulator *Arabis paniculata* Franch. *Environ. Exp. Bot.*, 66: 242-248.
- Zenk, M.H., 1996. Heavy metal detoxification in higher plants-a review. *Gene*, 179: 21-30.
- Zhuang, X., J. Chen, H. Shim and Z. Bai, 2007. New advances in plant growth-promoting rhizobacteria for bioremediation. *Environ. Int.*, 33: 406-413.